



**University of
Zurich^{UZH}**

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2019

Effects of intravenous infusion of E.coli lipopolysaccharide in early pregnant cows

Herzog, Kathrin ; Debertolis, Letizia ; Kastelic, John P ; Schmicke, Marion ; Ulbrich, Susanne E ; Bollwein, Heiner

Abstract: The objective was to characterize effects of *Escherichia coli* LPS (given iv) on corpus luteum (CL) and embryonic viability in early pregnant cattle. Eight non-lactating German Holstein cows were given 0.5 µg/kg LPS on 35 ± 3 d (mean ± SEM) of pregnancy, whereas seven heifers, 41 ± 6 d pregnant, were given 10 ml saline (Control group). Transrectal B-mode examinations of the CL were done at -1, 3, 6, 12, 24, 48, 72, and 96 h relative to treatment. Blood samples were collected at -1, 0.5, 1, 2, 3, 4, 6, 9, 12, 24, 48, 72, and 96 h. At 12 and 48 h, the CL was biopsied. None of the cows still in the experiment 10 d after LPS (n=7) had embryonic loss. In LPS-treated cows, luteal area decreased (from 4.1 to 3.1 cm²; P 0.05) within 6 h and until 48 h. Luteal blood flow decreased by 39% (P 0.05) within the first 6 h after LPS, but returned to pre-treatment values by 48 h. Plasma P4 decreased by 62% (P 0.05), reached a nadir (2.7±0.6 ng/mL) at 12 h after LPS and was not restored to pre-treatment (P 0.05). In luteal tissue, mRNAs for StAR and for FGF1 were lower (P 0.05) in LPS- than in saline-treated cattle at 12 h, with no difference between groups at 48 h. Levels of mRNAs for Casp3 and FGF2 were not different between groups (P>0.05) at 12 or 48 h after treatment. In conclusion, LPS transiently suppressed CL function, but did not induce embryonic mortality.

DOI: <https://doi.org/10.1530/REP-18-0174>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-158174>

Journal Article

Accepted Version

Originally published at:

Herzog, Kathrin; Debertolis, Letizia; Kastelic, John P; Schmicke, Marion; Ulbrich, Susanne E; Bollwein, Heiner (2019). Effects of intravenous infusion of E.coli lipopolysaccharide in early pregnant cows. *Reproduction*, 157(1):65-76.

DOI: <https://doi.org/10.1530/REP-18-0174>

Effects of intravenous infusion of *E.coli* lipopolysaccharide
in early pregnant cows

K. Herzog^{ab*}, L. Debertolis^{a*}, J. P. Kastelic^c, M. Schmicke^a, S. E. Ulbrich^d, and H.
Bollwein^e

- ^aClinic for Cattle, University of Veterinary Medicine Hannover, Germany
- ^bLower Saxony State Office for Consumer Protection and Food Safety,
Oldenburg, Germany (*Present Address*)
- ^cDepartment of Production Animal Health, University of Calgary, Canada
- ^dETH Zurich, Animal Physiology, Institute of Agricultural Sciences, Zurich, Switzerland
- ^eClinic of Reproductive Medicine, University of Zurich, Switzerland

*These authors contributed equally to this work.

*Correspondence and reprint requests:

Dr. Kathrin Herzog, Lower Saxony State Office for Consumer Protection and Food
Safety, Oldenburg, Germany

Phone: +49 (0)441 - 57026-288; Fax: +49 (0)441 - 57026-178;

E-mail: kathrin.herzog@laves.niedersachsen.de

Abstract

The objective was to characterize effects of *Escherichia coli* LPS (given iv) on corpus luteum (CL) and embryonic viability in early pregnant cattle. Eight non-lactating German Holstein cows were given 0.5 µg/kg LPS on 35 ± 3 d (mean ± SEM) of pregnancy, whereas seven heifers, 41 ± 6 d pregnant, were given 10 ml saline (Control group). Transrectal B-mode examinations of the CL were done at -1, 3, 6, 12, 24, 48, 72, and 96 h relative to treatment. Blood samples were collected at -1, 0.5, 1, 2, 3, 4, 6, 9, 12, 24, 48, 72, and 96 h. At 12 and 48 h, the CL was biopsied. None of the cows still in the experiment 10 d after LPS (n=7) had embryonic loss. In LPS-treated cows, luteal area decreased (from 4.1 to 3.1 cm²; $P \leq 0.05$) within 6 h and until 48 h. Luteal blood flow decreased by 39% ($P \leq 0.05$) within the first 6 h after LPS, but returned to pre-treatment values by 48 h. Plasma P4 decreased by 62% ($P \leq 0.05$), reached a nadir (2.7±0.6 ng/mL) at 12 h after LPS and was not restored to pre-treatment ($P \leq 0.05$). In luteal tissue, mRNAs for StAR and for FGF1 were lower ($P \leq 0.05$) in LPS- than in saline-treated cattle at 12 h, with no difference between groups at 48 h. Levels of mRNAs for Casp3 and FGF2 were not different between groups ($P > 0.05$) at 12 or 48 h after treatment. In conclusion, LPS transiently suppressed CL function, but did not induce embryonic mortality.

Key words: pregnancy; corpus luteum; endotoxin; gene expression; Color Doppler ultrasonography; progesterone; prostaglandin; cattle

47

Introduction

Embryonic mortality is of cardinal importance in bovine reproduction (Zavy 1994, Lucy 2001). Embryonic mortality is usually classified into two categories, based on the time of return to estrus. In cattle with early embryonic mortality, estrus occurs before day 24 post insemination (p.i.), compatible with embryonic death before day 16 p.i., whereas in late embryonic mortality, cattle return to estrus after day 24 p.i., apparently due to death of the embryo after day 16 p.i. (Kastelic & Ginther 1989, Humblot 2001, Santos *et al.* 2004a). Early embryonic mortality affects ~30 to 40% of all inseminated cattle, substantially reducing fertility (Thatcher *et al.* 1994, Silke *et al.* 2002, Diskin & Morris 2008). Late embryonic mortality is less frequent (5 to 20% of inseminated cattle) but causes greater economic losses per affected animal, as it substantially prolongs calving to conception intervals (Diskin & Sreenan 1980, Sreenan & Diskin 1983, Santos *et al.* 2004a).

Inflammatory processes during early pregnancy have been associated with embryonic mortality in cattle (Hansen *et al.* 2004), attributed to production of prostaglandin F₂alpha (PGF_{2α}) causing luteolysis and pregnancy loss (Manns *et al.* 1985, Stewart *et al.* 2003, Inskeep 2004). In cows given *Escherichia coli* lipopolysaccharides (*E. coli* LPS) during the first, second and third trimesters of pregnancy, 6 of 10 cows aborted (three, one and two in the first, second and third trimesters, respectively). Abortions were attributed to a prolonged release of PGF_{2α} (Giri *et al.* 1990) causing reductions in progesterone (P₄), especially during the first trimester (in the absence of an extraluteal source of P₄).

In a recent *in vivo* study, intravenous administration of *E.coli* LPS (model for

inflammation) temporarily decreased CL size and function in diestrus cows (Herzog *et al.* 2012). Luteal blood flow, plasma P₄ concentration and CL size decreased within 3, 9 and 24 h, respectively, after LPS. However, estrous cycle length was unchanged. Therefore, premature luteolysis did not occur.

The aim of the present study was to investigate effects of LPS exposure on the CL and on embryonic viability in early pregnant cattle. For this purpose, peripheral P₄ was used as main assessment of luteal secretory activity (Sartori *et al.* 2004, Lüttgenau *et al.* 2011). Furthermore, the relationship between endotoxin exposure and luteal function was evaluated at a subcellular level in biopsies of luteal tissue. Molecular parameters known to yield information regarding key cellular functions of the CL were selected (Niswender *et al.* 2000). Specifically, steroidogenic acute regulatory protein (StAR) is responsible for transportation of cholesterol to the inner mitochondrial membrane, a rate-limiting step in P₄ synthesis (Stocco & Clark 1996). Furthermore, Fibroblast Growth Factors (FGFs) participate in regulation of development and regression of the CL. In that regard, acidic and basic FGFs (FGF1 and FGF2, respectively) are increased during functional luteolysis (Doraiswamy *et al.* 1998, Neuvians *et al.* 2004); FGF2 is also involved in CL formation and controls P₄ secretion (Yamashita *et al.* 2008); and during luteolysis, Caspase-3 (Casp3) has a pivotal role in selective destruction of key structural and functional cellular proteins (Casciola-Rosen *et al.* 1996, Thornberry 1998).

Materials and methods

Cattle

This study was conducted (between April 2011 and May 2012) at the Clinic for Cattle, University of Veterinary Medicine, Hannover, Germany. The experimental protocol was reviewed and approved (Lower Saxony Federal State Office for Consumer Protection and Food Safety, 33.9-42502 – 04-11/0515) and the research was conducted in accordance with German legislation on animal welfare. Eight non-lactating German Holstein cows, clinically healthy and 35 ± 3 d (mean \pm SEM) after insemination, were used for the LPS trial. These cows were 3.7 ± 0.8 y old.

In addition, seven German Holstein heifers from the FLI (Friedrich Loeffler Institut – Federal Research Institute for Animal Health) in Mariensee, Germany were used as a control group (Saline group; $n=7$). They were 41 ± 6 d pregnant and 2.2 ± 0.3 y old, younger than the LPS-treated group ($P \leq 0.05$). All cattle were tethered in stalls and given *ad libitum* access to hay and water.

Study design

The LPS group were subjected to an Ovsynch protocol (Pursley *et al.* 1995), bred by timed artificial insemination, with pregnancy diagnosis (transrectal ultrasonography) done ~25 d after breeding. On day 35 ± 3 of pregnancy, a polyethylene catheter was inserted into a jugular vein. At 09.00 the following day, an LPS solution (0.5 μ g/kg body weight of *E. coli*, O55:B5; L2880, Sigma Aldrich, Steinheim am Albuch, Germany) in 10 ml sterile water (B. Braun, Melsungen) was given iv (over an interval of ~1 min). After administration of LPS, the jugular catheter was flushed with 25 ml 0.9% saline solution. Control heifers were bred following detection of estrus and given 10 ml of 0.9% saline solution instead of LPS on day 41 ± 6 of pregnancy.

Ultrasonography for assessment of the CL and pregnancy

Transrectal B-mode ultrasonographic examinations were done at -1, 3, 6, 12, 24, 48, 72, and 96 h relative to treatment (saline or LPS) to assess luteal size and luteal blood flow. A Logiq Book XP ultrasound scanner (General Electrics Medical Systems, Jiangsu, P.R. China), equipped with a 10.0-MHz linear-array transducer (General Electrics Yokogawa Medical Systems, Tokyo, Japan), was used. For luteal size, three cross-sectional images with maximal areas of the CL were recorded (using B-mode sonography) and luteal areas were measured offline (PixelFlux®, Version 1.0, Chameleon Software, Leipzig, Germany). If the CL had a cavity, the area of the cavity was measured and subtracted from total area (Kastelic *et al.* 1990). Means of the cross-sectional areas of the three images were calculated and used for statistical analyses. For luteal blood flow, power-flow Doppler was used for color blood flow mapping of the CL in various transverse sections, as described (Herzog *et al.* 2012).

To assess embryo/fetal viability, heartbeat was assessed with B-mode transrectal ultrasonography at 6, 12, 24, 48, 72, 96 h and at 10 d after infusion.

Blood samples and determination of plasma P_4 and prostaglandin metabolites concentrations

At 1.0 h before administration of LPS or saline solution, blood samples were collected to characterize concentrations of P_4 and PGFM. In addition, blood samples were collected (via the catheter) at 0.5, 1, 2, 3, 4, 6, 9, 12, 24, 48, 72, and 96 h after treatment.

Samples were stored on ice (0°- 4°C), centrifuged (3000 x g, 15 min at 4°C) within 30 min after collection and plasma removed stored at -20°C pending analyses.

Serum P₄ concentrations were determined with a commercial chemiluminescence immunoassay (Immulite®, Siemens Healthcare Diagnostics, Deerfield, IL, USA). The lower detection limit was 0.5 ng/ml, and intra- and inter-assay coefficients of variation were <10%. Plasma PGFM concentrations were determined with a competitive enzyme immunoassay (Mishra *et al.* 2003). The PGFM-horseradish peroxidase conjugate and antiserum were supplied by Prof. Meyer (Physiology Weihenstephan, Technische Universitaet Muenchen, Freising, Germany), whereas PGFM used for the standard curve was purchased from Sigma, Germany. The antiserum had minimal (<0.01%) cross reactions with PGE₂, PGEM, PGA₂, PGAM, and PGF_{2α} (Guvén & Ozsar 1993). The lowest detection limit for PGFM was 25 pg/ml. Intra- and inter-assay CVs were 3.5 and 11.4%, respectively.

Corpus luteum biopsy

Corpus luteum biopsies were done as described (Herzog *et al.* 2012) at 12 and 48 h after treatment. At each time of sampling, at least two tissue samples per animal were recovered and immediately placed in a sterile DNase- and RNase-free cryotube (Fa. Brand, Wertheim, Germany), frozen in liquid nitrogen, and stored at -80°C. This enabled repeated biopsy sampling from a single CL without impairing its function, as described (Tsai *et al.* 2001).

RNA extraction, cDNA synthesis and real-time qPCR

Total RNA was isolated from biopsy samples using QIAzol Lysis Reagent (QIAGEN Germany, Hilden), followed by two-step quantitative real-time RT-PCR (Pistek *et al.* 2013). A Bioanalyzer 2100 (Agilent Technologies, Waldbronn, Germany) and RNA 6000 Nano Kit (Agilent) were used to assess RNA Quality. Mean RIN values were 6.5 (range, 5.5-8.0). The qPCR was done with a CFX384™ Real-Time PCR Detection System (Bio-Rad, München, Germany) and PCR Mix SensiFast SYBR and Fluorescein Kit (Bioline, London, UK). Primers used are shown in Table 1. The cycle of quantification (Cq) was calculated after baseline subtracted curve fitting using the single threshold method (Bio-Rad CFX Manager V1.5.534.0511 software). Relative quantification of qPCR products was done as described (Livak & Schmittgen 2001). Geometric means of reference genes *H3F3A*, *UBK3*, *YWHAZ* were used for normalizing mRNA expression of target genes *FGF1*, *FGF2*, *StAR*, and *Casp3*, according to the BestKeeper method (Pfaffl *et al.* 2004).

Statistical analyses

For age, luteal size and blood flow, uterine blood flow parameters, and concentrations of P₄ and PGFM, residuals within groups were visually examined for normal distribution (PROC CHART) and analysed with a Shapiro-Wilk test (PROC UNIVARIATE). All statistical analyses were done with SAS® (version 9.1, Statistical Analysis Institute Inc., Cary, NC, USA).

Residuals within groups for the variable age did not differ significantly from a normal distribution. Hence, an independent two-sample Student's *t*-test (PROC TTEST) was used for comparison of the mean ages of the two groups.

Residuals within groups for variables LBF, *FGF1*, *FGF2*, *StAR* and *Casp3* also did not differ significantly from a normal distribution. Hence, these variables were submitted to a two-factorial variance analysis for repeated measurements (PROC GLM). The influence of various time points within groups was tested using repeated measurements variance analysis (PROC GLM). A Tukey's HSD test was used to locate differences between groups. A Student's *t*-test for paired samples (PROC MEANS) was used to compare measurement times within a group. Since variables P_4 , PGFM and LS did not have a normal distribution, nonparametric tests were used. Effects of measurement time on P_4 and LS were submitted to a Friedman's Test (PROC SORT) and differences between groups were located using Wilcoxon's rank sum test (PROC NPAR1WAY). Differences between time points within a group were tested using a Wilcoxon's signed rank test (PROC UNIVARIATE). For all statistical tests, a significance level of $P \leq 0.05$ was specified. Values for LBF, *FGF1*, *FGF2*, *StAR* and *Casp3* are expressed in text, figures and tables as mean \pm standard deviation (SD). For P_4 , PGFM and LS, median and median absolute deviation (MAD) were used.

Results

Clinical response

Treatment with LPS induced various toxemia-related clinical signs, including tachycardia, tachypnea, dyspnea, epiphora and mild diarrhea, as well as a slight increase of body temperature (maximal 39.7°C). However, all but one of the cattle were clinically normal by 12 h after treatment (Table 2). One cow was found dead 24 h after LPS treatment, with death attributed to cardiovascular failure. There were ~5 L of clotted

blood in the abdominal cavity, likely due to hemorrhage from the CL biopsy site, that certainly contributed to her demise. There was no pregnancy loss during or up to 10 d after LPS treatments (≤ 12 h: $n=8$; >12 h: $n=7$) and Control heifers did not have any clinical manifestations of toxemia.

Luteal area and luteal blood flow

Luteal size (LS) was 12% smaller ($P \leq 0.05$) at 6 h after LPS treatment, reached a nadir (3.1 ± 0.3 cm²) at 12 h, remained low until 24 h, and then returned to pre-treatment values by 48 h and thereafter remained unchanged (Fig. 1). For luteal blood flow (LBF), there were effects of treatment, time, and a treatment by time interaction ($P \leq 0.05$; Fig. 1). In LPS-treated cows, LBF decreased by 39% ($P \leq 0.05$) at 6 h, reached a nadir (0.9 ± 0.2 cm²) at 12 h, and then rebounded to pre-treatment values by 48 h (Fig. 1). In Control heifers, LS and LBF remained relatively stable.

Plasma P_4 and PGFM concentrations

For plasma P_4 concentrations, there were effects of time ($P \leq 0.05$) and a treatment by time interaction ($P \leq 0.05$; Fig. 2). Plasma P_4 concentrations increased from 7.1 ± 0.7 to 10.3 ± 2.7 ng/ml (median \pm MAD; $P \leq 0.05$) within the first 30 min after administration. Thereafter, P_4 decreased between consecutive measurements ($P \leq 0.05$) beginning 4 h after infusion, reached a nadir (2.7 ± 0.6 ng/ml) 12 h after infusion, increased again between 24 and 48 h ($P \leq 0.05$), but was not restored to pre-treatment values. In Control heifers, P_4 concentrations did not vary significantly between consecutive measurements during the first 12 h ($P > 0.05$). Progesterone concentrations decreased ($P \leq 0.05$) between

12 and 24 h after saline infusion, increased again between 24 and 48 h ($P \leq 0.05$) and thereafter remained constant. Plasma P_4 concentrations were higher in LPS-treated than control cattle within the first 30 min after treatment ($P \leq 0.05$), lower ($P \leq 0.05$) in LPS-treated cattle between 9 and 12 h after treatment and higher ($P \leq 0.05$) in LPS-treated cattle at 48 h.

For PGFM, there were effects of time, and a treatment by time interaction ($P \leq 0.5$; Fig. 2). Plasma PGFM concentrations increased almost by an order of magnitude within the first 30 min after infusion of LPS, but had returned to baseline by 4 h and remained there. In Control heifers, PGFM did not change significantly over time. Concentrations of PGFM were higher ($P > 0.05$) in LPS versus Control cattle for the first 4 h after infusion.

Gene expression

In cows given LPS, transcript abundance of mRNA encoding *Casp3* was higher at 12 versus 24 h after treatment ($P \leq 0.5$, Fig. 3), whereas expression of *StAR* mRNA was lower ($P \leq 0.5$, Fig. 3) at 12 versus 48 h. There was no difference within the LPS group for *FGF2* ($P > 0.05$), whereas *FGF1* was lower ($P \leq 0.05$, Fig. 3) at 12 versus 48 h after LPS infusion. Within the Control group, expression of mRNA did not vary between time points ($P > 0.05$) for any parameter. Levels of mRNA encoding *StAR* and *FGF1* were lower in LPS- than in saline-treated cattle at 12 h after treatment ($P \leq 0.05$; Fig. 3) but not at 48 h. There were no differences between treatment groups for *Casp3* or *FGF2* ($P > 0.05$).

Discussion

In early-pregnant cows in the present study, iv administration of LPS endotoxin temporarily suppressed CL function and structure. Luteal size, LBF, and P_4 decreased within 12 h after LPS treatment, but complete luteal regression did not occur. Similarly, in a previous study, administration of LPS to diestrous cows reduced LS, LBF and P_4 during the first 24 h after treatment (Herzog *et al.* 2012), but did not cause luteal regression.

In the present study, all fetuses still in the experiment at the time ($n=7$) had a heart beat 10 d after treatment. Progesterone is essential for maintenance of pregnancy in cattle (Estergreen *et al.* 1967). In the first trimester of gestation, removal of the CL causes abortion. Progesterone of placental origin can support pregnancy at ~150 to 250 d of gestation in the absence of a CL. However, removal of the CL after 250 d results in calving in 1 or 2 d (Stabenfeldt *et al.* 1970, Mann & Lamming 1999, Taverne 2001, Breukelman *et al.* 2005). Furthermore, pregnancy is maintained by exogenous P_4 in cattle in absence of endogenous P_4 , e.g. with the CL removed by ovariectomy (Kesler 1997) or administration of luteolytic doses of $PGF_{2\alpha}$ (Ferguson *et al.* 2014). In particular, based on a recent report (Ferguson *et al.* 2014) the sooner P_4 supplementation starts after $PGF_{2\alpha}$, the more likely the pregnancy will be maintained, with a positive outcome most likely if exogenous P_4 is given within 12 h after $PGF_{2\alpha}$. Furthermore, those authors observed that P_4 decreased fastest between 12 and 24 h after $PGF_{2\alpha}$, and was below 2 ng/ml by 30 h post- $PGF_{2\alpha}$. In this study, P_4 in LPS-treated cows was also suppressed for more than 12 h in a similar pattern, but consistently exceeded 2 ng/ml. Luteal regression and declines in plasma P_4 to < 2 ng/ml after administration of exogenous $PGF_{2\alpha}$ are well established (Lauderdale 1975, Sioan 1977, Wright & Kiracofe 1988). We concluded that

the LPS-induced release of endogenous $\text{PGF}_{2\alpha}$ impaired luteolytic function but did not cause luteolysis, thereby reducing P_4 production, but not getting systemic P_4 concentrations low enough to terminate pregnancy.

Administering viable bacteria or prolonged infusions of LPS activate the immunological cascade, disrupt neuroendocrine activity (mainly by blocking or delaying the LH surge), and can shorten the estrous cycle in postpartum cattle (Williams *et al.* 2008). Also, studies using mastitis as a model of inflammation instead of experimentally LPS administration have implicated infectious diseases as causing embryonic loss (Soto *et al.* 2003, Santos *et al.* 2004b). However, in the present study, LPS was administered over a short interval (~1 min). Therefore, the insult was short-lived. Consequently, only one PGFM peak was detected; although it clearly suppressed P_4 , it did not cause luteolysis and the CL subsequently recovered. In heifers, intrauterine infusions of exogenous $\text{PGF}_{2\alpha}$ caused a PGFM pattern with several pulses, similar to spontaneous luteolysis (Ginther *et al.* 2009). That the single PGFM peak detected in the present study did not induce luteolysis was further evidence that sequential PGF pulses are crucial for complete luteolysis in cattle (Ginther *et al.* 2009, Ginther *et al.* 2010).

In third-trimester cows ($n=3$, 190 to 200 d of gestation) given LPS (0.5 μg Salmonella typhimurium LPS/kg) intravenously, the LPS was rapidly cleared and there were no indications that it crossed the placenta. However, all three cows aborted within 10 d after LPS treatment (Foley *et al.* 1993). In the present study, none of the cows aborted. Perhaps the third-trimester fetus is more susceptible to LPS than cattle at ~6 wk of pregnancy. Also, differences in source and dose of LPS could have contributed to disparate outcomes. Effects of two doses of LPS (1.0 or 2.5 μg *E. coli* LPS/kg) in cows

at various stages of pregnancy have been reported (Giri *et al.* 1990). In that study, LPS was infused over 6 h (which better represented exposure to LPS in natural infection), in contrast to the single bolus injection given in the present study. Furthermore, magnitudes of changes in P_4 , PGFM, cortisol and other metabolites, as well as severity of clinical signs, were all dose-dependent. Although the type of LPS used was the same as in the present work, both doses were higher. Intensity of systemic responses to LPS in cattle is dose-dependent, and can considerably vary from cow to cow (Jacobsen *et al.* 2005).

Progesterone concentrations after LPS treatment in the present study were similar to those in cyclic cows (Herzog *et al.* 2012). In both experiments, P_4 peaked 30 min after LPS exposure. This was attributed to an acute release of P_4 from the adrenal gland, as in the study by Battaglia *et al.* (1997), with a nadir in P_4 after 12 h. However, P_4 concentrations in cyclic LPS treated cows remained decreased for up to 48 h after LPS exposure (Herzog *et al.* 2012), whereas the difference between early pregnant LPS and control cattle had disappeared by 24 h after LPS administration. Progesterone concentrations also recovered much faster in pregnant cows of the present study than in cyclic cows (Herzog *et al.* 2012). Furthermore, P_4 concentrations were higher 48 h after treatment in LPS-treated cows compared to controls. Similarly, for LBF, there was no difference between LPS and saline-treated pregnant cattle in LBF 24 h after LPS administration, and LBF was higher in LPS-treated cows 48 and 72 h after treatment, consistent with a robust association between P_4 and LBF, as reported (Herzog *et al.* 2010). Changes in luteal size caused by LPS infusion were less prominent in pregnant than in cyclic cows. Luteal size sharply decreased in pregnant cows, reached a nadir at

12 h and returned to initial values by 48 h after infusion, whereas in cyclic cows, the nadir occurred after 24 h and LS never regained pre-treatment baseline (Herzog *et al.* 2012).

Luteal gene expression confirmed that the CL of LPS-treated cows in the present study underwent partial luteolysis. As in cyclic cows (Herzog *et al.* 2012), *StAR* was reduced 12 h after LPS administration in pregnant cows. Already at 48 h, however, there were no differences between LPS and control cattle in expression of *StAR* mRNA. Unfortunately, in the previous experiment, expression of mRNA in cyclic cows was not assessed 48 h after infusion. Regardless, it is noteworthy that plasma P₄ concentrations differed between groups for only a short interval.

There were no significant differences between LPS-treated versus Control cows for mRNA expression of *Casp3*. Conversely, in cyclic cows (Herzog *et al.* 2012) expression of this protein was increased 12 h after administration in LPS-treated cows compared to controls, in accordance with decreased luteal size. Perhaps *Casp3* has an important role in apoptosis and/or subsequent regeneration of cycling, but not in CL of pregnant cows, where it may have a lesser effect. Observations in laboratory animals support this assertion. In that regard, *Casp3* can be dispensable or nonspecific in mice (Kuida *et al.* 1996, Jänicke *et al.* 1998). Furthermore, in postpartum rats, *Casp3* only regulated regression of the CL generated after parturition, but had no apparent role in luteolysis of a CL formed during pregnancy (Takiguchi *et al.* 2004). Further investigation is needed to determine the relevance of these findings for the bovine CL. In addition, we evaluated *Casp3* only at mRNA level. Assessing the presence of the *Casp3* protein would provide

further information (Johnson & Bridgham 2000, Fenwick & Hurst 2002); therefore, it should be considered in future studies.

There were significant differences in expression of *FGF1* but not *FGF2* after LPS exposure. Similarly, in a previous study, *FGF1* was reduced in cyclic LPS treated cows 12 h after exposure (Lüttgenau *et al.* 2016). Both *FGF1* and *FGF2* are regarded as cellular “survival factors” (Renaud *et al.* 1994), conferring protection against apoptosis. They promote angiogenesis in the developing CL, as well as support pre-existing vasculature in the mature and gravid CL (Zheng *et al.* 1993, Schams 1994). Therefore, reductions in *FGF1* and a corresponding decrease in LBF at 12 h after LPS infusion were expected. In cyclic cows, both *FGF1* and *FGF2* were increased at 12 h after induction of luteolysis (Neuvians *et al.* 2004). In contrast, in the present study, *FGF2* expression did not change over time and *FGF1* expression was lower 12 h after LPS infusion. Furthermore, *FGF2* was increased in diestrous cows given LPS (Lüttgenau *et al.* 2016). Since the cited experiments were done in cyclic cattle, apparent differences may have been due to pregnancy status. There may be luteoprotective mechanisms in the CL of pregnant cattle that make them less susceptible to endogenous PGF, consistent with the ‘rebound’ in CL function in LPS-treated cows in the present study.

The LPS-treated cows underwent an OvSynch protocol and timed artificial insemination, whereas heifers in the control group were inseminated after estrus detection. This difference was due to organizational reasons, but is potentially relevant. Timed artificial insemination is a potential risk factor for late embryonic loss, although direct evidence is missing (Santos *et al.* 2004a). However, rather than luteal or uterine events, follicular development seems to play a relevant part in the mechanism by which

368 timed artificial insemination affects pregnancy rate (Jordan *et al.* 2009, Bollwein *et al.*
369 2010). Fortunately, none of the LPS-cows in our study underwent embryonic loss.

370 Due to organizational reasons, there was also a difference in parity number and
371 age between LPS- and Control group (primiparous cows vs. nulliparous heifers).
372 However, none of the parameters measured (LS, LBF, P₄ and PGFM) differed between
373 the two groups before LPS infusion. Furthermore, the minimal reaction to the NaCl
374 infusion of the heifers in this study resembled that of cycling cows (Herzog *et al.* 2012).
375 In addition, the Control group was mainly designed to exclude any influence of stressors
376 (e.g. manipulations related to treatment, sonography and other examinations), since, in
377 general, stress impairs reproductive function (Sheldon *et al.* 2014, Endo *et al.* 2017). In
378 this regard, there is evidence that heifers might be even less stress-resistant than cows.
379 When cattle are repeatedly exposed to stressors such as painless restraint, transport
380 and manipulation, their reaction to those stimuli progressively decreases. This applies to
381 behavioral alterations as well as to hormonal response, e.g. cortisol level (Boissy &
382 Bouissou 1988, Lay Jr *et al.* 1996, Grandin 1997, Rushen *et al.* 2001). Because of their
383 younger age, nulliparous heifers are obviously less accustomed to human contact and
384 manipulations than cows. Thus, they are likely more stress-sensitive than older cows
385 (Lay Jr *et al.* 1996). An inverse relation between parity number and intensity of reaction
386 to stressors has been reported in other species, including gestating sows (Zhang *et al.*
387 2017).

388 One cow died between 12 and 24 h after LPS infusion. At necropsy, there were
389 approximately 5 L of clotted blood in the abdomen, apparently derived from luteal biopsy
390 sites. The ultrasound-guided technique described by (Tsai *et al.* 2001) allows repeated

biopsies from a CL, without apparent effect on its function. To our knowledge, this is the first case reporting hemorrhage from the biopsy site after using this technique. It is known from human medicine that leukopenia and thrombocytopenia develop in immunocompetent patients during septic shock (Stephan *et al.* 1997). In a healthy cow, the loss of 5 liters of blood would not cause death (Radostits *et al.* 2000). However, in an animal that was already compromised, this may have acted in a synergistic manner with acute phase symptoms, resulting in death.

In summary, administration of LPS to early pregnant cows caused a temporary depression of luteal function, but all pregnancies were preserved. Furthermore, the impact on CL was less pronounced than in a similar study involving diestrous cows. In particular, analysis of molecular parameters suggested differences in luteolytic mechanisms between the CL of the cycle and the CL of pregnancy, consistent with luteoprotective changes in the latter.

Declaration of interest

The authors declare that there is no conflict of interest that could be perceived as prejudicing the impartiality of the research reported.

Funding

This research did not receive any specific grant from any funding agency in the public, commercial, or not-for-profit sectors.

Acknowledgments

414 The authors thank the Friedrich-Löffler-Institute in Mariensee for generously providing
415 access to pregnant heifers.

416

References

- Battaglia DF, Bowen JM, Krasa HB, Thrun LA, Viguie C & Karsch FJ** 1997 Endotoxin inhibits the reproductive neuroendocrine axis while stimulating adrenal steroids: A simultaneous view from hypophyseal portal and peripheral blood. *Endocrinology* **138** 4273-4281.
- Boissy A & Bouissou M-F** 1988 Effects of early handling on heifers' subsequent reactivity to humans and to unfamiliar situations. *Applied Animal Behaviour Science* **20** 259-273.
- Bollwein H, Prost D, Ulbrich SE, Niemann H & Honnens A** 2010 Effects of a shortened preovulatory follicular phase on genital blood flow and endometrial hormone receptor concentrations in Holstein-Friesian cows. *Theriogenology* **73** 242-249.
- Breukelman SP, Szenci O, Beckers J-F, Kindahl H, Mulder EJ, Jonker FH, van der Weijden B, Revy D, Pogany K & Sulon J** 2005 Ultrasonographic appearance of the conceptus, fetal heart rate and profiles of pregnancy-associated glycoproteins (PAG) and prostaglandin F 2 α -metabolite (PGF 2 α -metabolite) after induction of fetal death with aglepristone during early gestation in cattle. *Theriogenology* **64** 917-933.
- Casciola-Rosen L, Nicholson DW, Chong T, Rowan KR, Thornberry NA, Miller DK & Rosen A** 1996 Apopain/CPP32 cleaves proteins that are essential for cellular repair: a fundamental principle of apoptotic death. *The Journal of Experimental Medicine* **183** 1957-1964.
- Diskin MG & Morris DG** 2008 Embryonic and early foetal losses in cattle and other ruminants. *Reproduction in Domestic Animals* **43 Suppl 2** 260-267.
- Diskin MG & Sreenan JM** 1980 Fertilization and embryonic mortality rates in beef heifers after artificial insemination. *Journal of Reproduction and Fertility* **59** 463-468.
- Doraiswamy V, Knutson DL, Grazul-Bilska AT, Redmer DA & Reynolds LP** 1998 Fibroblast growth factor receptor (FGFR)-1 and -2 in the ovine corpus luteum throughout the estrous cycle. *Growth Factors* **16** 125-135.

- Endo N, Kuroki R & Tanaka T 2017 Comparison of productive and reproductive performance and hair cortisol levels between Brown Swiss cross-bred and Holstein cows housed in the same barn. *Animal Science Journal* **88** 1506-1512.
- Estergreen VL, Frost OL, Gomes WR, Erb RE & Bullard JF 1967 Effect of ovariectomy on pregnancy maintenance and parturition in dairy cows. *Journal of Dairy Science* **50** 1293-1295.
- Fenwick M & Hurst P 2002 Immunohistochemical localization of active caspase-3 in the mouse ovary: growth and atresia of small follicles. *Reproduction* **124** 659-665.
- Ferguson EC, Kesler DJ & Godke RA 2014 Maintenance of pregnancy in beef cattle after a luteolytic dose of prostaglandin F2 α . *Veterinary Medicine and Animal Sciences* **2** 1-4.
- Foley GL, Schlafer DH, Elsasser TH & Mitchell M 1993 Endotoxemia in pregnant cows: Comparisons of maternal and fetal effects utilizing the chronically catheterized fetus. *Theriogenology* **39** 739-762.
- Ginther OJ, Araujo RR, Palh  o MP, Rodrigues BL & Beg MA 2009 Necessity of sequential pulses of prostaglandin F2 α for complete physiologic luteolysis in cattle. *Biology of Reproduction* **80** 641-648.
- Ginther OJ, Shrestha HK, Fuenzalida MJ, Shahiduzzaman AKM, Hannan MA & Beg MA 2010 Intrapulse temporality between pulses of a metabolite of prostaglandin F2 α and circulating concentrations of progesterone before, during, and after spontaneous luteolysis in heifers. *Theriogenology* **74** 1179-1186.
- Giri SN, Emau P, Cullor JS, Stabenfeldt GH, Bruss ML, Bondurant RH & Osburn BI 1990 Effects of endotoxin infusion on circulating levels of eicosanoids, progesterone, cortisol, glucose and lactic acid, and abortion in pregnant cows. *Veterinary Microbiology* **21** 211-231.
- Grandin T 1997 Assessment of stress during handling and transport. *Journal of animal science* **75** 249-257.
- Guven B & Ozsar S 1993 Set up an enzymimmunoassay for PGFM and its release during the estrous cycle in the Angora goat. *Proceedings of the world congress on animal production, Edmonton, Canada* p. 289.

- Hansen PJ, Soto P & Natzke RP 2004 Mastitis and fertility in cattle - possible involvement of inflammation or immune activation in embryonic mortality. *American Journal of Reproductive Immunology* **51** 294-301.
- Herzog K, Brockhan-Ludemann M, Kaske M, Beindorff N, Paul V, Niemann H & Bollwein H 2010 Luteal blood flow is a more appropriate indicator for luteal function during the bovine estrous cycle than luteal size. *Theriogenology* **73** 691-697.
- Herzog K, Struve K, Kastelic JP, Piechotta M, Ulbrich SE, Pfarrer C, Shirasuna K, Shimizu T, Miyamoto A & Bollwein H 2012 Escherichia coli lipopolysaccharide administration transiently suppresses luteal structure and function in diestrous cows. *Reproduction* **144** 467-476.
- Humblot P 2001 Use of pregnancy specific proteins and progesterone assays to monitor pregnancy and determine the timing, frequencies and sources of embryonic mortality in ruminants. *Theriogenology* **56** 1417-1433.
- Inskeep EK 2004 Preovulatory, postovulatory, and postmaternal recognition effects of concentrations of progesterone on embryonic survival in the cow. *Journal of animal science* **82** (E-Suppl) E24-39.
- Jacobsen S, Toelboell T & Andersen PH 2005 Dose dependency and individual variability in selected clinical, haematological and blood biochemical responses after systemic lipopolysaccharide challenge in cattle. *Veterinary Research* **36** 167-178.
- Jänicke RU, Sprengart ML, Wati MR & Porter AG 1998 Caspase-3 is required for DNA fragmentation and morphological changes associated with apoptosis. *Journal of Biological Chemistry* **273** 9357-9360.
- Johnson A & Bridgham J 2000 Caspase-3 and -6 expression and enzyme activity in hen granulosa cells. *Biology of Reproduction* **62** 589-598.
- Jordan A, Herzog K, Ulbrich SE, Beindorff N, Honnens A, Krueger L, Miyamoto A & Bollwein H 2009 Genital blood flow and endometrial gene expression during the preovulatory period after prostaglandin F₂ α -induced luteolysis in different luteal phases in cows. *Journal of Reproduction and Development* **55** 309-315.

- Kastelic JP & Ginther OJ 1989 Fate of conceptus and corpus luteum after induced embryonic loss in heifers. *Journal of the American Veterinary Medical Association* **194** 922-928.
- Kastelic JP, Pierson RA & Ginther OJ 1990 Ultrasonic morphology of corpora lutea and central luteal cavities during the estrous cycle and early pregnancy in heifers. *Theriogenology* **34** 487-498.
- Kesler DJ 1997 Norgestomet implants maintain pregnancy in ovariectomized heifers. *Theriogenology* **48** 89-98.
- Kuida K, Zheng TS, Na S, Kuan C-Y, Yang D, Karasuyama H, Rakic P & Flavell RA 1996 Decreased apoptosis in the brain and premature lethality in CPP32-deficient mice. *Nature* **384** 368-372.
- Lauderdale J. 1975. The use of prostaglandins in cattle. In *Annales de Biologie Animale Biochimie Biophysique* **2**, pp. 419-425.
- Lay Jr D, Friend T, Randel R, Jenkins O, Neuendorff D, Kapp G & Bushong D 1996 Adrenocorticotrophic hormone dose response and some physiological effects of transporation on pregnant Brahman cattle. *Journal of animal science (USA)*.
- Livak KJ & Schmittgen TD 2001 Analysis of relative gene expression data using real-time quantitative PCR and the 2- $\Delta\Delta$ CT method. *Methods* **25** 402-408.
- Lucy MC 2001 Reproductive loss in high-producing dairy cattle: where will it end? *Journal of Dairy Science* **84** 1277-1293.
- Lüttgenau J, Beindorff N, Ulbrich SE, Kastelic JP & Bollwein H 2011 Low plasma progesterone concentrations are accompanied by reduced luteal blood flow and increased size of the dominant follicle in dairy cows. *Theriogenology* **76** 12-22.
- Lüttgenau J, Herzog K, Strüve K, Latter S, Boos A, Bruckmaier RM, Bollwein H & Kowalewski MP 2016 LPS-mediated effects and spatio-temporal expression of TLR2 and TLR4 in the bovine corpus luteum. *Reproduction* **151** 391-399.
- Mann GE & Lamming GE 1999 The influence of progesterone during early pregnancy in cattle. *Reproduction in Domestic Animals* **34** 269-274.
- Manns JG, Nkuuhe JR & Bristol F 1985 Prostaglandin concentrations in uterine fluid of cows with pyometra. *Canadian Journal of Comparative Medicine* **49** 436-438.

- Mishra DP, Meyer HH & Prakash BS 2003 Validation of a sensitive enzymeimmunoassay for 13,14-dihydro-15-keto-PGF₂α in buffalo plasma and its application for reproductive health status monitoring. *Animal Reproduction Science* **78** 33-46.
- Neuvians TP, Berisha B & Schams D 2004 Vascular endothelial growth factor (VEGF) and fibroblast growth factor (FGF) expression during induced luteolysis in the bovine corpus luteum. *Mol Reprod Dev* **67** 389-395.
- Niswender GD, Juengel JL, Silva PJ, Rollyson MK & McIntush EW 2000 Mechanisms controlling the function and life span of the corpus luteum. *Physiol Rev* **80** 1-29.
- Pfaffl MW, Tichopad A, Prgomet C & Neuvians TP 2004 Determination of stable housekeeping genes, differentially regulated target genes and sample integrity: BestKeeper--Excel-based tool using pair-wise correlations. *Biotechnol Lett* **26** 509-515.
- Pistek VL, Furst RW, Kliem H, Bauersachs S, Meyer HH & Ulbrich SE 2013 HOXA10 mRNA expression and promoter DNA methylation in female pig offspring after in utero estradiol-17β exposure. *Journal of Steroid Biochemistry and Molecular Biology* **138** 435-444.
- Pursley JR, Mee MO & Wiltbank MC 1995 Synchronization of ovulation in dairy cows using PGF₂α and GnRH. *Theriogenology* **44** 915-923.
- Radostits OM, Gay CC, Blood DC & Hinchcliff KW 2000. Diseases of the blood and blood-forming organs. In *Veterinary Medicine. A textbook of the Diseases of Cattle, Sheep, Pigs, Goats and Horses*.
- Renaud F, Oliver L, Desset S, Tassin J, Romquin N, Courtois Y & Laurent M 1994 Up-regulation of aFGF expression in quiescent cells is related to cell survival. *Journal of Cellular Physiology* **158** 435-443.
- Rushen J, Munksgaard L, Marnet P & DePassillé A 2001 Human contact and the effects of acute stress on cows at milking. *Applied Animal Behaviour Science* **73** 1-14.
- Santos JE, Thatcher WW, Chebel RC, Cerri RL & Galvao KN 2004a The effect of embryonic death rates in cattle on the efficacy of estrus synchronization programs. *Animal Reproduction Science* **82-83** 513-535.

- Santos JEP, Cerri RLA, Ballou MA, Higginbotham GE & Kirk JH 2004b** Effect of timing of first clinical mastitis occurrence on lactational and reproductive performance of Holstein dairy cows. *Animal Reproduction Science* **80** 31-45.
- Sartori R, Haughian JM, Shaver RD, Rosa GJM & Wiltbank MC 2004** Comparison of ovarian function and circulating steroids in estrous cycles of Holstein heifers and lactating cows. *Journal of Dairy Science* **87** 905-920.
- Schams D 1994** Growth factors in milk. *Endocrine Regulations* **28** 3-8.
- Sheldon IM, Cronin JG, Healey GD, Gabler C, Heuwieser W, Streyll D, Bromfield J, Miyamoto A, Fergani C & Dobson H 2014** Innate immunity and inflammation of the bovine female reproductive tract in health and disease. *Reproduction* **148** R41-R51.
- Silke V, Diskin MG, Kenny DA, Boland MP, Dillon P, Mee JF & Sreenan JM 2002** Extent, pattern and factors associated with late embryonic loss in dairy cows. *Animal Reproduction Science* **71** 1-12.
- Sioan C 1977** Induced abortion in cattle. *Theriogenology* **7** 43-51.
- Soto P, Natzke RP & Hansen PJ 2003** Identification of possible mediators of embryonic mortality caused by mastitis: actions of lipopolysaccharide, prostaglandin F₂α, and the nitric oxide generator, sodium nitroprusside dihydrate, on oocyte maturation and embryonic development in cattle. *American Journal of Reproductive Immunology* **50** 263-272.
- Sreenan JM & Diskin MG 1983** Early embryonic mortality in the cow: its relationship with progesterone concentration. *Veterinary record* **112** 517-521.
- Stabenfeldt GH, Osburn BI & Ewing LL 1970** Peripheral plasma progesterone levels in the cow during pregnancy and parturition. *American Journal of Physiology* **218** 571-575.
- Stephan F, Thiolliere B, Verdy E & Tulliez M 1997** Role of hemophagocytic histiocytosis in the etiology of thrombocytopenia in patients with sepsis syndrome or septic shock. *Clinical infectious diseases* **25** 1159-1164.

- 593 **Stewart AB, Inskeep EK, Townsend EC & Dailey RA** 2003 Effects of Gram-positive
594 bacterial pathogens in ewes: peptidoglycan as a potential mediator of interruption
595 of early pregnancy. *Reproduction* **125** 295-299.
- 596 **Stocco DM & Clark BJ** 1996 Regulation of the acute production of steroids in
597 steroidogenic cells. *Endocr Rev* **17** 221-244.
- 598 **Takiguchi S, Sugino N, Esato K, Karube-Harada A, Sakata A, Nakamura Y, Ishikawa H**
599 **& Kato H** 2004 Differential regulation of apoptosis in the corpus luteum of
600 pregnancy and newly formed corpus luteum after parturition in rats. *Biology of*
601 *Reproduction* **70** 313-318.
- 602 **Taverne MAM** 2001 Corpus luteum function and parturition in cattle and pigs. *ARCHIV*
603 *FUR TIERZUCHT* **44** 37-50.
- 604 **Thatcher WW, Staple CR, Danet-Desnoyers G, Oldick B & Schmitt EP** 1994 Embryo
605 health and mortality in sheep and cattle. *Journal of animal science* **72; Suppl 3**
606 16-30.
- 607 **Thornberry NA** 1998 Caspases: key mediators of apoptosis. *Chemistry & Biology* **5** R97-
608 103.
- 609 **Tsai SJ, Kot K, Ginther OJ & Wiltbank MC** 2001 Temporal gene expression in bovine
610 corpora lutea after treatment with PGF2alpha based on serial biopsies in vivo.
611 *Reproduction* **121** 905-913.
- 612 **Williams EJ, Herath S, England GWC, Dobson H, Bryant CE & Sheldon IM** 2008 Effect
613 of Escheria coli infection of the bovine uterus from the whole animal to the cell.
614 *Animal* **2** 1153-1157.
- 615 **Wright JM & Kiracofe GH** 1988 Short estrous cycles and associated progesterone in
616 serum of beef heifers aborted at various stages of gestation. *Theriogenology* **29**
617 497-504.
- 618 **Yamashita H, Kamada D, Shirasuna K, Matsui M, Shimizu T, Kida K, Berisha B, Schams**
619 **D & Miyamoto A** 2008 Effect of local neutralization of basic fibroblast growth
620 factor or vascular endothelial growth factor by a specific antibody on the
621 development of the corpus luteum in the cow. *Mol Reprod Dev* **75** 1449-1456.
- 622 **Zavy MT** 1994. Embryonic mortality in cattle. In *Embryonic mortality in domestic species*.

623 **Zhang M-y, Li X, Zhang X-h, Liu H-g, Li J-h & Bao J** 2017 Effects of confinement
624 duration and parity on behavioural responses and the degree of psychological
625 fear in pregnant sows. *Applied Animal Behaviour Science* **193** 21-28.

626 **Zheng J, Redmer DA & Reynolds LP** 1993 Vascular development and heparin-binding
627 growth factors in the bovine corpus luteum at several stages of the estrous cycle.
628 *Biology of Reproduction* **49** 1177-1189.
629

Figure legends

Fig. 1: Median \pm MAD of luteal tissue area (Panel A) and mean \pm SEM of luteal blood flow (Panel B) in early pregnant cattle treated with *E. coli* lipopolysaccharide (LPS; ■; -1 to 12 h: n=8, 24 to 96 h: n=7) or Saline (○; n=7), respectively.

*Values differed between *LPS-* and *Saline-treated* cattle ($P \leq 0.05$).

^{a,b}Within *LPS-treated* cattle, values without a common superscript differed ($P \leq 0.05$).

Fig. 2: Median \pm MAD of serum progesterone (Panel A) and PGFM concentration (Panel B) in early pregnant cattle treated with *E. coli* lipopolysaccharide (LPS; ■; -1 to 12 h: n=8, 24 to 96 h: n=7) or Saline (○; n=7), respectively.



*Values differed between *LPS-* and *Saline-treated* cattle ($P \leq 0.05$).


^{a,b,A,B}Within *LPS-treated* cattle, values without a common superscript differed ($P \leq 0.05$).

Fig. 3: Mean \pm SEM levels of mRNA for StAR (Panel A), Casp3 (Panel B), FGF1 (Panel C) and FGF2 (Panel D) 12 and 48 h after infusion of *E. coli* lipopolysaccharide (LPS; dark bars) or *Saline* (NaCl; grey bars) in early pregnant cattle.


* Values differed between *LPS-* and *Saline-treated* cattle within a sampling period (i.e. 12 or 48 h), ($P \leq 0.05$).


^{a,b}Within *LPS-treated* cattle, values without a common superscript differed ($P \leq 0.05$)





B





Gene	Gene symbol	reference [accession no.]	forward primer [5'...3']	reverse primer [3'...5']	PCR-product [bp]	AT [°C]	FA [°C]	MP [°C]
<i>polyubiquitin</i>	<i>UBQ3</i>	NM_174133	AGATCCAGGATAAGGAAGGCAT	GCTCCACCTCCAGGGTGAT	198	60	83	88
<i>histone</i>	<i>H3F3A</i>	NM_174133	AGATCCAGGATAAGGAAGGCAT	GCTCCACCTCCAGGGTGAT	233	60	80	87
<i>tyrosine 3-monooxygenase</i>	<i>YWHAZ</i>	XM_001927228	AGGCTGAGCGATATGATGAC	GACCCTCCAAGATGACCTAC	141	60	81	87
<i>fibroblast growth factor 1 (acidic)</i>	<i>FGF1</i>	NM_174055	GCTGAAGGAGAAACCAGCAC	GTTTTCTCCAACCTTTCCA	108	60	75	82
<i>fibroblast growth factor 2 (basic)</i>	<i>FGF2</i>	NM_174056	TCAAAGGAGTGTGTGTGAAC	CAGGGCCACATACCAACTG	288	60	75	82
<i>toll-like receptor 2</i>	<i>TLR-2</i>	NM_174197	CCATGTGGAGAGGGTGTT	GGGGACACAAAACAGCACTT	140	60	81	85
<i>toll-like receptor 4</i>	<i>TLR-4</i>	NM_174198	GACCCTTGCGTACAGTTGT	GGTCCAGCATCTTGGTTGAT	103	60	83	88
<i>caspase-3</i>	<i>CASP-3</i>	NM_214131	AAC CTC CGT GGA TTC AAA ATC	TTC AGG RTA ATC CAT TTT GTA AC	114	60	75	81
<i>steroidogenic acute regulatory</i>	<i>StAR</i>	XR_083945	GGATTAACCAGGTTCCGGCG	CTCTCCTTCTCCAGCCCTC	157	60	84	89

1
2

Table 2. Clinical symptoms of each LPS-treated cow. Abbreviations: T= rectal temperature; HR= heart rate; RR= respiratory rate; ESV= episcleral vessels

	1	2	3	4	5	6	7	8
Time in relation to LPS (hh:mm)	-1:00	-1:00	-1:00	-1:00	-1:00	-1:00	-1:00	-1:00
	T 38.6°C HR 76/min RR 24/min No abnormal clinical signs	T 38.4°C HR 56/min RR 36/min No abnormal clinical signs	T 38.7°C HR 56/min RR 24/min No abnormal clinical signs	T 38.5°C HR 64/min RR 24/min No abnormal clinical signs	T 38.5°C HR 72/min RR 32/min No abnormal clinical signs	T 38.6°C HR 60/min RR 24/min No abnormal clinical signs	T 38.8°C HR 72/min RR 40/min No abnormal clinical signs	T 38.7°C HR 56/min RR 24/min No abnormal clinical signs
	Tachycardia RR to 60/min, eupneic tachypnea Standing, bright, sudden interruption of feed intake	Tachycardia RR to a maximum of 90/min Mild to severe dyspnea, coughing, grunting, stridor Standing, bright Polyuria Epiphora	Standing, bright Tachycardia RR to 52/min Moderate dyspnea, coughing, mild stridor	RR to 36/min Mild dyspnea, coughing Standing, bright	Tachycardia RR to 54/min Mild dyspnea, dilated nostrils, coughing Standing, bright, sudden interruption of feed intake	Standing, bright Tachycardia RR to 56/min Moderate dyspnea, coughing	RR 40/min Standing, bright, nervous	Tachycardia RR 90/min Moderate dyspnea, coughing Standing, becomes less responsive, sudden interruption of feed intake
	0:15 to 0:30	Tachycardia RR to a maximum of 104/min Moderate dyspnea with stridor, sporadic grunting, coughing Standing, becomes less responsive, head down	Tachycardia RR 90/min, irregular Moderate to severe dyspnea, frequent and deep grunting Standing, eupneic position (head down and forward, abducted front limbs), anxious, less responsive Eructation 1-2 times per minute	Standing Tachycardia RR to 52/min Moderate dyspnea, coughing, stridor	Tachycardia RR 60 to 80/min Moderate dyspnea with stridor, coughing Standing, becomes less responsive, nervous, head down Ptyalism Epiphora	Tachycardia RR 54 to 104/min Moderate to severe dyspnea with coughing, mouth breathing Standing, eupneic position (holds head forward), nervous, becomes less responsive Ptyalism Mild diarrhea	Standing, bright Tachycardia RR to 100/min Moderate to severe dyspnea with stridor, coughing, mouth breathing	Tachycardia RR to 54/min Moderate dyspnea, coughing Standing, bright, anxious
	0:30 to 1:00	Tachycardia RR to 80/min Moderate dyspnea with stridor, frequent grunting, sporadic coughing Unsteadiness, alternates weight bearing on left vs right rear limbs Ptyalism	Tachycardia RR 80/min Severe dyspnea with stridor, grunting, coughing Unsteady, alternates weight between rear limbs Ptyalism, epiphora and seromucous nasal discharge	Standing, anxious, unresponsive, holds head down Tachycardia RR to a maximum of 60/min Severe dyspnea with stridor, grunting, coughing Ptyalism Moderate diarrhea	Tachycardia RR 80/min Moderate dyspnea with stridor, grunting, coughing Standing, anxious, unresponsive, ears cold Ptyalism Epiphora and injected ESV Mild diarrhea	Tachycardia RR to 92/min Severe dyspnea with grunting, coughing Standing, nervous, unsteadiness, alternates weight on rear limbs, unresponsive, head down Ptyalism Mild diarrhea	Tachycardia RR to 68/min Moderate dyspnea with stridor, grunting Standing, less responsive, unsteady, alternated weight on rear limbs Ptyalism Moderate diarrhea	Tachycardia RR 90/min Moderate to severe dyspnea with dilated nostrils, stridor, grunting, coughing, epistaxis Standing, eupneic position (head down and forward, abducted front limbs), anxious, unsteady, alternates weight between rear limbs, becomes unresponsive
	1:00 to 2:00	T 39.4°C Tachycardia RR 52 to 80/min Moderate dyspnea with stridor Frequent grunting Anxious, drinks repeatedly, Unresponsive Ptyalism	T 38.4°C Tachycardia RR from 24 to 48/min Severe dyspnea with stridor, frequent grunting Anxious, searches repeatedly for water but does not drink, Unresponsive, up and down Ptyalism, epiphora and seromucous nasal discharge Ears cold (only apical) Injected ESV Mild diarrhea	T 38.8°C Tachycardia RR 44 to 48/min Moderate dyspnea with stridor, coughing Standing, is calm, but still unresponsive Injected ESV Mild diarrhea	T 38.5°C Tachycardia RR 64 to 72/min Severe dyspnea with stridor, grunting, coughing, mouth breathing Up and down, anxious, unresponsive Epiphora and injected ESV Mild diarrhea	T 39.4°C Tachycardia RR 78 to 88/min Moderate dyspnea, stridor, grunting Standing, eupneic position (holds head forward), anxious, unresponsive, alternates weight on rear limbs Ptyalism Injected ESV Tremors	T 38.6°C Tachycardia RR to 64/min Moderate dyspnea, grunting Mostly standing, unresponsive Bristled hair Injected ESV	T = 39.3°C RR 60/min Mild dyspnea, coughing Up and down anxious, unsteadiness, alternates weight between rear limbs, unresponsive, tremors Injected ESV Moderate diarrhea Ptyalism Injected ESV
	2:00 to 3:00	T 39.3°C Tachycardia RR 56 to 60/min Moderate dyspnea with stridor, frequent grunting and coughing Up and down Calm, but unresponsive Ptyalism Injected ESV	T 39.0°C RR 32/min to 24/min Severe to moderate dyspnea, stridor, grunting sporadically Mostly recumbent, frequently shifted between lying on left vs right Unresponsive Injected ESV Moderate diarrhea Other signs tend to disappear	T 38.8°C Tachycardia RR 44/min to 36/min Moderate to mild dyspnea with stridor Standing, is calm, but still unresponsive, drinks repeatedly Ptyalism disappears Injected ESV Mild diarrhea	T 38.5°C Tachycardia RR 64/min to 54/min Moderate dyspnea with grunting Up and down, calm, but unresponsive Ptyalism Epiphora and injected ESV Mild diarrhea	T 39.0°C Tachycardia RR to 80/min Moderate dyspnea, grunting Standing, calm, but unresponsive, drinks repeatedly Injected ESV	T 38.5°C RR to 64/min Mild dyspnea, grunting Mostly standing, unresponsive Injected ESV	T 39.7°C RR to 60/min Mild dyspnea, stridor, grunting, coughing Recumbent, nervous, unresponsive Moderate diarrhea Ptyalism Injected ESV Tremors
	3:00 to 6:00	T 38.6°C Tachycardia gradually decreases RR slowed from 56 to 36/min (3:00 to 5:00) Moderate to mild dyspnea with gurgling and stridor; grunting and cough disappear gradually 5:00 becomes calm and responsive Ptyalism Injected ESV 5:00 to 6:00 polyuria	T 38.5°C RR 27/min Mild dyspnea tends to decrease, slight stridor, grunting sporadically Mostly standing, is calm, but still unresponsive Injected to normal ESV Mild diarrhea	T 38.4°C 3:00 No tachycardia RR 36/min to 20/min Mild dyspnea tends to disappear Up and down, becomes calm and responsive Tremors Injected ESV Mild diarrhea	T 38.2°C Tachycardia gradually decreases RR 40 to 48/min Moderate to severe to mild dyspnea with stridor, grunting, coughing, gradually decreasing Up and down, head down, unresponsive Tremors Ptyalism Injected ESV Mild diarrhea	T 37.5°C (4:17) to 38.0°C (5:00) No tachycardia (4:17) RR to 56/min Mild dyspnea, coughing, grunting Up and down, still unresponsive Mild dehydration Tremors Injected ESV	T 39.3°C RR to 56/min Mild dyspnea, grunting Mostly standing, becomes calm and responsive (4:53) Injected ESV	T 39.4°C (3:06) to 39.0 (4:08) RR to 44/min Mostly recumbent, becomes calm and responsive All signs progressively disappear
	6:00 to 12:00	T 38.4°C RR 36/min Mild dyspnea and stridor gradually disappeared 6:00, feed and water intake restored 7:50 all signs disappeared	T 38.5°C 8:51 HR and RR in normal range, no abnormal clinical signs	T 38.6°C Mostly recumbent, calm, bright RR 24/min 07:25 no abnormal clinical signs	T 38.3°C RR 40/min No dyspnea Mostly recumbent 7:15 becomes calm and responsive, feed and water intake restored All other signs gradually disappear 9:45 no abnormal clinical signs	T 38.7°C RR 44 to 58/min Mild dyspnea only when recumbent, cough Mostly recumbent, unresponsive, feed and water intake restored Injected ESV All other signs disappear (11:17)	T 39.3°C RR to 40/min Mild dyspnea, then decreased Mostly recumbent, bright, feed and water intake restored (6:00) 9:03 no abnormal clinical signs	T 38.8°C Mostly recumbent, bright 6:06 HR and RR in normal range, no abnormal clinical signs
	12:00 to 24:00 (end)	12:00 T, HR and RR in normal range, no abnormal clinical signs	12:00 T, HR and RR in normal range, no abnormal clinical signs 21:00 Found dead (apparently died between 15:00 and 20:00)	12:00 T, HR and RR in normal range, no abnormal clinical signs	12:00 T, HR and RR in normal range, no abnormal clinical signs	12:00 T, HR and RR in normal range, no abnormal clinical signs	12:00 T, HR and RR in normal range, no abnormal clinical signs	12:00 T, HR and RR in normal range, no abnormal clinical signs